

		Active Mobilization															
		Thumb*			Index Finger*			Middle Finger*			Ring Finger*			Little Finger*			
Loss of Amplitude of Movements	Muscle Strength (4 or 5/5)	IP	MP	CM	DIP	PIP	MP	DIP	PIP	MP	DIP	PIP	MP	DIP	PIP	MP	
		Normal limits	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Decrease in amplitude of movement, functional position maintained	6	6	6	1.5	1.5	0.75	2	2	1	1	1	0.5	1.5	1.5	0.75
		Total ankylosis in functional position	12	10	10	4	4	2	6	6	3	3	3	1.5	4	4	2
		Total ankylosis in incomplete or faulty position	20	12	12	8	4	3	10	6	4	5	3	2	8	4	3
Amputation		20	12	12	8	4	3	10	6	4	5	3	2	8	4	3	

When the amputation of a phalanx is partial, the score used is the one indicated for the joint closest to the site of the amputation.
 In the case of the distal phalanx, no score is given if more than 50% of the normal length of the phalanx is preserved.

Muscle Weakness (3/5 or less)	20	12	12	8	4	3	10	6	4	5	3	2	8	4	3
--------------------------------------	----	----	----	---	---	---	----	---	---	---	---	---	---	---	---

Total of Table B = _____ Points

- * IP: Interphalagial
- PIP: Proximal Interphalagial
- DIP: Distal Interphalagial
- MP: Metacarpo-phalagial
- CM: Carpo-metacarpal